

Recovery Assessment of Open-heart Cardiac Surgery Patients Using Heart Rate Variability Parameters

Seyedsadra Miri¹, Sabina Lähteenmäki^{1,2}, Lassi Tuomisto^{1,2}, Heidi Mahrberg², Antti Vehkaoja¹, Jari Laurikka^{1,2}, Jari Viik¹

¹Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland

²Tampere University Hospital Heart Center, Tampere, Finland

Abstract

Surgical operation causes short-term stress changing the balance of the autonomic nervous system (ANS). The ANS activity can be assessed through heart rate variability (HRV). This work aims to evaluate the recovery process of open-heart cardiac surgery patients under two post-surgery physiotherapy interventions using HRV parameters. Ten-minute ECG recordings were performed on 17 open-heart cardiac surgery patients pre-operatively (PREOP) and during three consecutive post-operative days (POD1, POD2 and POD3). The recovery process was promoted using two different physiotherapy techniques: positive expiratory pressure (PEP, N = 9) and inspiratory training (IMT, N = 8). Common HRV parameters (SDNN, LF power and SD₁) were calculated for each patient. Statistical analyses inside and between the PEP and IMT groups were performed by dependent and independent parametric and non-parametric tests.

The results show significant differences between HRV parameters pre- and post-operatively. Cross-subject averages of SDNN, LF and SD₁ in PREOP and POD1 for PEP group recovery intervention were 36.46 and 18.07 ms, 636.08 and 42.84 ms², 21.06 and 5.89 ms, respectively and for IMT group they were 32.05 and 19.66 ms, 283.31 and 73.18 ms², 13.69 and 8.98 ms, respectively. There was a reduction of all three investigated HRV parameters in the POD1 for both groups (e.g., for SDNN: $p = 0.02$ and $p = 0.042$ for PEP and IMT groups, respectively). LF power average values increased from 42.84 to 159.46 ms² and 73.18 to 119.02 ms² for the PEP and IMT groups, respectively, between POD1 to POD3 indicating that patients were eventually starting to recover from the surgery. Statistically significant difference between PEP and IMT interventions were not discovered (e.g., $p = 0.54$ for SDNN, PREOP vs. POD3). Based on the results, all studied HRV parameters are potential indicators of the short-term recovery after cardiac surgery.

1. Introduction

Autonomic nervous system (ANS) acts as a control system for the regulation of internal processes of body and is influenced by environmental and physiological factors. Its two branches, sympathetic (SNS) and parasympathetic (PNS) branch behave in an opposing way to maintain cardiovascular homeostasis. While SNS is responsible for the heart rate (HR) and blood pressure (BP) increase, PNS regulation leads to the decrease of HR and BP [1]. Cardiac surgeries as well as other major operations under general anesthesia can affect the status of ANS due to the short-term applied physiological stress [2]. Thus, non-invasively monitoring of surgical patients is of high importance in order to assess the recovery procedure. Heart rate variability (HRV) monitoring is a non-invasive monitoring technique which has been widely utilized as an assessment method for cardiac autonomic modulation [3]. It has been shown that HRV can be a useful tool for the predication of cardiac post-surgery complications [4] [5] [6], assessing arrhythmias [7], and evaluating the effects of post-surgery therapeutic intervention [8].

In order to boost the recovery period of surgery patients, different physiotherapy intervention techniques such as positive expiratory pressure (PEP) or inspiratory muscle training (IMT) can be utilized. PEP technique is based on applying a pressure to create resistance against breathing which can increase lung volume and reduce hyperinflation [9] whereas IMT method exercises diaphragm-based muscles with inspiratory function [10].

In this study we assessed the recovery procedure of the open-heart cardiac surgery patients under two different therapeutic interventions and see how surgical interventions and their effects on ANS can be reflected through HRV indices and how different therapeutic interventions can boost the recovery process.

2. Materials and methods

Ten-minute ECG recordings were performed for 17 open-heart cardiac surgery patients at Tampere University

Hospital Heart Center in Tampere, Finland during 2013–2016. All patients participated in the study voluntarily and gave their informed consent in written. The study was approved by the regional Ethics Committee of Tampere University Hospital area, Finland.

For each patient the data was collected during one pre-operative day (PREOP) and three post-operative days (POD1, POD2 and POD3). Two different therapeutic interventions were utilized during the recovery process; positive expiratory pressure (PEP group, N = 9) and inspiratory muscle training (IMT group, N = 8).

2.1. Measurement procedure

The ECG was recorded with a small custom-designed impedance pneumography device and sampled at 256 Hz frequency [11]. Bipolar ECG measurement having disposable electrodes in the arms close to the armpit was used. ECG data was stored during ten-minute period of tidal breathing.

2.2 ECG signal processing

Baseline wandering and high frequency noise caused by powerline interference were removed from the raw ECG recordings using low-pass and median high-pass filters. This made the signals prepared for R-peak extraction. R-peaks were localized and extracted using the method proposed by Zhengzhong *et al.* [12]. The HRV tachograms were then formed to be further analyzed. Atrial and ventricular ectopic beats were detected and removed using time series artifact correction algorithm proposed by Lipponen *et al.* [13]. Matlab R2019b was used for the analysis.

2.3. HRV parameters and statistical analysis

Common HRV parameters, SDNN, LF and SD₁, were calculated for 10-minute segments. Statistical analysis was performed between surgery days and between intervention groups. Normality distribution was assessed by Shapiro-Wilk test and independent Student’s t-test or Mann-Whitney U test, and Paired Student’s t-test or Wilcoxon Signed Rank Test were used to determine significance level. The significance level was considered 0.05.

3. Result

Table 1 shows cross-subject averages of SDNN, LF power and SD₁ values of PEP and IMT intervention groups pre-operatively and during three post-operative days.

Table 1. Cross-subject averages of SDNN, LF power and SD₁ pre- and post-operatively for PEP and IMT intervention groups.

	PREOP		POD1		POD2		POD3	
	PEP	IMT	PEP	IMT	PEP	IMT	PEP	IMT
SDNN [ms]	36,4	32	18	19	16,8	18	18,2	19,3
LF power [ms ²]	636	283	42,8	73	56,1	177	159	119
SD ₁ [ms]	21	13,	5,9	8,9	6,8	10	10,6	9,4

The difference between baseline values of intervention groups for SDNN, LF power and SD₁ were measured, and p-values were 0.603, 0.888 and 0.196, respectively thus no statistically significant difference was found in the investigated HRV parameter values between the groups pre-operatively.

The significance of differences between pre-operative day and three post-operative days for different HRV parameters is provided in Table 2. The last column of the table compares the intervention methods during the process of recovery (PREOP vs POD3) using parametric and non-parametric statistical tests.

Table 2. Significance levels (p-values) of baseline compared to post-operative days and comparison of interventions during the recovery period. P-values less than 0.05 were considered significant and are marked by asterisks (*).

	PREOP vs POD1		PREOP vs POD2		PREOP vs POD3		PREOP vs POD3 (PEP & IMT)
	PEP	IMT	PEP	IMT	PEP	IMT	
SDNN [ms]	0,02 *	0,04 *	0,02 *	0,02 *	0,07	0,05	0,540
LF power [ms ²]	0,06	0,02 *	0,11	0,03 *	0,16	0,01 *	0,888
SD ₁ [ms]	0,01 *	0,06	0,01 *	0,05	0,07	0,06	0,409

Figure 1 shows the SDNN, LF power and SD₁ for each subject. It can be seen that few subjects have different trends of recovery procedure.

4. Discussion and conclusion

The achieved results clearly show that there is a significant reduction of SDNN, LF power and SD₁ on the first post-operative day for both intervention groups which is a result of surgical stress caused by the open-heart cardiac surgery and its effects on ANS. It has been shown by Brown *et al.* that the full recovery of ANS can take approximately 6 to 12 weeks after the surgery [14]. The rate of the recovery has large individual variation, but the process is only done during 3-day timeframe (POD1 to

POD3) that was the duration of the recovery monitoring in the present study. Therefore, HRV parameter values cannot be expected to recover back to the baseline level

within this timeframe. However, it can be seen that most HRV parameters start to slightly increase during the recovery period (POD1 to POD3).




Figure 1. SDNN, LF power and SD₁ values for each subject pre- and post-operatively for both intervention groups. Statistically significant difference between pre- and post-operative days is determined by asterisks (p-value < 0.05).

Due to the low number of participants in this study (N = 17), the average baseline values differ between the two intervention groups. However, not statistically significant difference was found between HRV parameters pre-operatively; making it possible to further compare the intervention groups on post-operative days. The recovery procedure of intervention groups was compared, and no significant statistical difference was found between PEP and IMT intervention groups for any of the HRV parameters.

In conclusion, it was found that HRV parameters can be utilized as a tool for monitoring of the recovery of open-heart cardiac surgery patients. It was also concluded that different therapeutic interventions did not have any effects of recovery procedure. However, in order to develop practical solutions for recovery monitoring, larger studies with more participants, longer monitoring period and various surgery groups shall be conducted.

References

[1] Z. M. Jenkins, N. Eikelis, A. Phillipou, D. J. Castle, H. E. Wilding and E. A. Lambert, "Autonomic Nervous System Function in Anorexia Nervosa: A Systematic Review,"

Frontiers in neuroscience, vol. 15, pp. 682208-682208, 28 June 2021.

[2] N. Herring, M. Kalla and D. J. Paterson, "The autonomic nervous system and cardiac arrhythmias: current concepts and emerging therapies," *Nature reviews cardiology*, vol. 16 (12), pp. 760-760, 2019.

[3] A. M. Catai, C. M. Pastre, M. F. d. Godoy, E. d. Silva, A. C. d. M. Takahashi and L. C. M. Vanderlei, "Heart rate variability: are you using it properly? Standardisation checklist of procedures," *Revista brasileira de fisioterapia*, vol. 24 (2), pp. 91-102, March 2020.

[4] A. Nenna, M. Lusini, C. Spadaccio, F. Nappi, S. M. Greco, R. Barbato, E. Covino and M. Chello, "Heart rate variability : a new tool to predict complications in adult cardiac surgery," *geriatric cardiology*, vol. 14 (11), pp. 662-668, 2017.

[5] N. Lakusic, D. Mahovic, P. Kruzliak, J. Cerkez Habek, M. Novak and D. S. M. S. Cerovec, "Changes in Heart Rate Variability after Coronary Artery Bypass Grafting and Clinical Importance of These Findings," *BioMed research international*, vol. 2015, pp. 680515-7, 20 May 2015.

[6] C. B. F. Pantoni, R. G. Mendes, L. Di Thommazo-Luporini, R. P. Simões, O. Amaral-Neto, R. Arena, S. Guizzilini, W. J. Gomes, A. M. Catai and A. Borghi-Silva, "Recovery of linear and nonlinear heart rate dynamics after

coronary artery bypass grafting surgery," *Clinical physiology and functional imaging*, vol. 34 (6), pp. 449-456, November 2014.

- [7] S. Guzzetti, E. Borroni, P. E. Garbelli, E. Ceriani, P. Della Bella, N. Montano, C. Cogliati, V. K. Somers, A. Mallani and A. Porta, "Symbolic dynamics of heart rate variability : A probe to investigate cardiac autonomic modulation," *Circulation*, vol. 112 (4), pp. 465-470, 2005.
- [8] N. Y. Tamburus, R. F. L. Paula, V. C. Kunz, M. C. César, M. A. Moreno and E. da Silva, "Interval training based on ventilatory anaerobic threshold increases cardiac vagal modulation and decreases high-sensitivity c-reactive protein: randomized clinical trial in coronary artery disease," *Revista brasileira de fisioterapia*, vol. 19 (6), pp. 441-450, 2015.
- [9] M. Fagevik Olsén, L. Lannefors and E. Westerdahl, "Positive expiratory pressure – Common clinical applications and physiological effects," *Respiratory medicine*, vol. 109 (3), pp. 297-307, 2014.
- [10] J. Wu, L. Kuang and L. Fu, "Effects of inspiratory muscle training in chronic heart failure patients: A systematic review and meta-analysis," *Congenital heart disease*, vol. 13 (2), pp. 194-202, 2018.
- [11] T. Vuorela, V.-P. Seppa, J. Vanhala and J. Hyttinen, "Design and Implementation of a Portable Long-Term Physiological Signal Recorder," *IEEE transactions on information technology in biomedicine*, vol. 14 (3), pp. 718-725, 2010.
- [12] G. Zhengzhong, K. Fanxue and Z. Xu, "Accurate and Rapid QRS Detection for Intelligent ECG Monitor," in *2011 Third International Conference on Measuring Technology and Mechatronics Automation*, 2011.
- [13] J. A. Lipponen and M. P. Tarvainen, "A robust algorithm for heart rate variability time series artefact correction using novel beat classification," *Journal of medical engineering & technology*, vol. 43 (3), pp. 173-181, 03 April 2019.
- [14] C. A. Brown, L. A. Wolfe, S. Hains, G. Ropchan and J. Parlow, "Heart rate variability following coronary artery bypass graft surgery as a function of recovery time, posture, and exercise," *Canadian journal of physiology and pharmacology*, vol. 82 (7), pp. 457-464, 01 July 2004.

Address for correspondence:

Seyedsadra Miri
Faculty of Medicine and Health Technology, Tampere
University, Arvo Ylpön Katu 34 (33520 Tampere), P.O.
Box 100, FI-33014, Tampere, Finland
seyedsadra.miri@tuni.fi