

A Causal Discovery Approach to Streamline Ionic Currents Selection to Improve Drug-Induced TdP Risk Assessment

Safaa Al-Ali¹, Jordi Llopis-Lorente², Maria Teresa Mora², Maxime Sermesant¹, Beatriz Trenor², Irene Balelli¹

¹ Centre INRIA d'Université Côte d'Azur, EPIONE team, Sophia Antipolis, France

² Centro de Investigación e Innovación en Bioingeniería (Ci2B), Universitat Politècnica de València, Valencia, Spain

Abstract

Causality is of paramount importance in bio-medical data analysis: assessing the causal relationships between the observed variables allows to improve our understanding of the tackled condition and better support decision-making. Torsade de Pointes (TdP) is a serious drug-induced cardiac side effect which can lead to sudden death. TdP is related to abnormal repolarizations in single cells, and the minimum set of ion channels needed to correctly assess TdP risk is still an open question. In this work, we propose to apply the causal discovery method ICA-Linear Non-Gaussian Acyclic Model (ICA-LiNGAM) to uncover the relations across the 7 ion channels identified by the Comprehensive in vitro Pro-arrhythmia Assay (CiPA) initiative as potentially related to the induction of TdP: I_{Kr} , I_{Na} , I_{NaL} , I_{CaL} , I_{K1} , I_{Ks} and I_{to} . We consider 109 drugs of known torsadogenic risk listed by CredibleMeds. We identify I_{Kr} , I_{NaL} and I_{CaL} as the ones that directly affect TdP-risk assessment, and suggest that I_{Na} perturbations could potentially have a high impact on pro-arrhythmic risk induction. Our causality-based results were further confirmed by independently performing binary drug risk classification, which shows that the combination of the 3 selected ionic currents maximizes the classification accuracy and specificity, outperforming state-of-the-art approaches based on alternative ion channel combinations.

1. Introduction

Identifying causal links between variables of importance takes a leading position in facilitating stable inference and making rational decisions in several areas, by overcoming the bias induced through standard correlation-based statistical and machine-learning approaches. For instance, in medicine, understanding the causal relationship between risk factors and disease can help develop effective prevention and treatment strategies and prioritize the available in-

formation to improve risk assessment tools.

Causal discovery is a branch of causal research, whose aim is to learn the cause-effect relationships of observational variables, typically through causal graphs. Several methods have been proposed to learn such graphs, for instance by exploring assumptions on the independent causal function representing the relation between each variable (or node) and its parents' nodes in the graph, or the independent variable-specific noise term [1]. Among them, the linear non-Gaussian acyclic model (LiNGAM) [2] assumes that the causal relations between variables are simply linear and their noise term is non-Gaussian to enforce the identification of the causal directions. An extension of the LiNGAM algorithm, ICA-LiNGAM further assumes that the observed variables are not independent due to the presence of unobserved latent variables [3]: independent component analysis (ICA, [4]) is used to separate the observed variables into their independent components, hence LiNGAM is applied.

Drug-induced Torsade de Pointes (TdP) is one of the most frightening drug side effects since it can trigger ventricular fibrillation and ultimately lead to sudden death. TdP is closely associated with abnormal repolarization, hence by a prolongation of the QT interval. The human ether-à-go-go-related gene (hERG) is responsible for the rapid component of the delayed rectifier current (I_{Kr}) which is one of the major repolarizing currents in the heart [5]. Proarrhythmic risk assessment of drugs is traditionally based on the evaluation of the hERG channel and the measure of the delayed ventricular repolarization on the electrocardiogram (ECG), i.e. the QT interval prolongation [6]. This method can accurately classify high TdP-risk drugs through a single analytical assessment considering a unique ion channel and exclusively focusing on ventricular repolarization [7]. However, it has been observed (eg [8]) that I_{Kr} blocking may be not sufficient to assess drug-induced TdP risk and is prone to produce false positives. Consequently, for a more reliable drug safety assessment,

recent studies have been proposed to incorporate multiple ion channels, since drug-altered currents may interact and cause disrupted cardiac electrophysiology [9–11]. Several ion channels’ combinations have been retained in the literature as the most relevant to assess drug-induced TdP risk, such as I_{Kr} , I_{CaL} , I_{Na} [9, 12] or I_{Kr} , I_{CaL} , I_{NaL} , and I_{Ks} [10].

In this work, we propose to use ICA-LiNGAM for upstream ion channels’ selection, to improve TdP risk assessment. We introduce the causal discovery algorithm ICA-LiNGAM and the considered drug data set. In Section 3, we present our results: the uncovered relationships between the ion channels and the binary drugs’ label, and their straightforward applicability for the classification task. Section 4 provides our conclusions and discussion of future directions for the presented work.

2. Materials and method

2.1. Drugs data-set

A total of 109 drugs from CredibleMeds¹ with known torsadogenic risk (37 with known, 14 with possible, 13 with conditional, and 45 with no proven risk) were used in this study. More details about the data set are available in [10] and the supplementary material herein. We consider a binary classification task, labeling drugs of confirmed or possible TdP-risk as unsafe, and drugs with conditionally or no proven TdP risk as safe.

For every drug, we use two pharmacological data: the IC_{50} for each of the seven ionic currents addressed by the CiPA initiative, and the effective free therapeutic plasma concentration (EFTPC), defined as the drug concentration in the plasma required to produce the desired therapeutic effect in the body. We consider the ion channels blocked fraction, here denoted by Bf_{ion} :

$$Bf_{ion} := \left[1 + \left(\frac{IC_{50}Ion}{EFTPC} \right)^h \right]^{-1}, \quad (1)$$

where h denotes the Hill coefficient, the number of drug molecules assumed to be sufficient to block an ion channel.

2.2. ICA-LiNGAM algorithm

ICA-LiNGAM [3], summarized in Algorithm 1, is a function-based causal discovery algorithm [13], which extends the Linear Non-Gaussian Additive Model (LiNGAM) [2]. LiNGAM assumes that the causal relations between observed variables can be represented by a non-Gaussian linear acyclic model, i.e. a directed acyclic graph (DAG) [14] where the causal functions between each variable and its parents’ nodes are linear, and the error term

follows a centered non-Gaussian distribution with non-zero variance. Let $\mathbf{X} = \{x_i, i = 1, \dots, n\}$ be the set of observed variables: we can encode the DAG structure through its adjacency matrix $\mathbf{B} = \{b_{ij}\}$, where b_{ij} represents the strength of the connection between the variables x_i and x_j . The matrix \mathbf{B} could be permuted to be strictly lower triangular, in accordance with the acyclicity assumption. Finally, denoting by Pa_i the set of parents nodes of x_i (i.e. the nodes who causally precede x_i), the generating process for variable $x_i \in \mathbf{X}$ writes:

$$x_i = \sum_{x_j \in Pa_i} b_{ij}x_j + e_i, \quad (2)$$

where $e_i \in \mathbf{E}$ are the error terms. We can rewrite Equation (2) in a matrix form,

$$\mathbf{X} = (\mathbb{I} - \mathbf{B})^{-1}\mathbf{E}, \quad (3)$$

\mathbb{I} being the identity matrix. The ICA-LiNGAM method considered here is based on the additional assumption that observed variables may be dependent due to the presence of unobserved latent variables [3]. Indeed, Equation (3) defines the independent component analysis model since the noise terms e_i are independent and non-Gaussian. ICA [4] is used to estimate the mixing matrix $\mathbf{A} = (\mathbb{I} - \mathbf{B})^{-1}$. Further, a set of permutations and scaling of the obtained independent parameters are performed to obtain the causal order k and the adjacency matrix as well.

Algorithm 1 ICA-LiNGAM

Input: Data matrix \mathbf{X}

Output: $\tilde{\mathbf{B}}$, estimation of \mathbf{B} (Equation (3))

1. Given \mathbf{X} , use ICA to obtain the decomposition $\mathbf{X} = \mathbf{A}\mathbf{S}$, where \mathbf{S} contains the independent components. Define $\mathbf{W} := \mathbf{A}^{-1}$.
 2. Compute the matrix $\tilde{\mathbf{W}}$, the unique permutation of \mathbf{W} without any leading diagonal zero.
 3. Compute $\tilde{\mathbf{W}}'$, which corresponds to $\tilde{\mathbf{W}}$ after dividing each row by its corresponding diagonal element.
 4. Compute an estimation $\hat{\mathbf{B}}$ of $\mathbf{B} := \mathbb{I} - \tilde{\mathbf{W}}'$.
 5. Estimate the causal order for each variable $x_i \in \mathbf{X}$ by finding the permutation matrix $\tilde{\mathbf{P}}$ of $\hat{\mathbf{B}}$ such that $\tilde{\mathbf{B}} := \tilde{\mathbf{P}}\hat{\mathbf{B}}\tilde{\mathbf{P}}^T$ is as close as possible to a strictly lower triangular structure ($\sum_{i \leq j} \tilde{\mathbf{B}}_{ij}^2$).
 6. Return the permuted matrix $\tilde{\mathbf{B}}$.
-

3. Results

In Figure 1 we show the causal graph obtained by applying ICA-LiNGAM, using the python package `gcasle`, to the ion channels blocked fractions, for the seven ionic currents identified by the CiPA initiative: I_{Kr} , I_{Na} , I_{NaL} ,

¹See CredibleMeds (accessed March 9, 2020)

I_{CaL} , I_{K1} , I_{Ks} and I_{to} . We introduce an extra *Label* node to investigate the relations of the blockade parameters with the known pro-arrhythmic risk. We used 5-fold cross-validation over the 109 drugs, and the final graph represented in Figure 1 accounts for the occurrences of causal arrows identified in each fold. A directed causal arrow from variable A to variable B indicates that changes in the parent node A can directly cause changes in the children node B . One can see that the I_{Kr} channel is systemically identified as a direct cause leading to the drug label, which is directly caused as well by I_{CaL} and I_{NaL} channels. Even though I_{Na} channel is not affecting the node label directly, it still plays an important role since it is causally related to I_{Kr} , I_{CaL} , and I_{NaL} channels. Indeed, the obtained DAG suggests that if a drug has an impact on I_{Na} blockade, it will also have an impact on its descendant ion channels blockade, I_{Kr} , I_{CaL} , I_{NaL} .




Figure 1: Causal graph generated by ICA-LiNGAM, through 5-fold cross-validation. The arrow thickness represents the number of occurrences of the causal arrow over the 5 splits.

Next, we perform an independent binary classification of the 109 drugs used for this study. In particular, we used two classical machine learning classifiers: Random-Forest and K-Nearest-Neighbors, whose final prescriptions are combined through a majority voting classifier [15]. A 5-fold cross-validation was performed as well for the classification task. In Figure 2 we present the accuracy and specificity scores for the ion channels combinations revealed by the obtained causal graph (Figure 1), starting from the most stably identified parent of the label node, I_{Kr} , up to its farthest ancestors. The highest accuracy levels are achieved by the two combinations (I_{Kr} , I_{CaL} , I_{NaL}) and (I_{Kr} , I_{CaL} , I_{NaL} , I_{Na}). For the ion-combination (I_{Kr} , I_{CaL} , I_{NaL}) we reach a mean accuracy level of 93.59%, whereas the combination (I_{Kr} , I_{CaL} , I_{NaL} , I_{Na}) allows to achieve a mean accuracy of 92.68%.

Furthermore, our causality-based classification shows a high mean specificity, reaching 94.70% for (I_{Kr} , I_{CaL} , I_{NaL}). There is no further improvement beyond this combination, indicating that the causal graph has been effective in iden-




Figure 2: Accuracy and specificity of drugs’ safety classification based on the combinations of ionic currents suggested by the DAG in Figure 1, using the voting classifier with Random-Forest and K-Nearest-Neighbors. The white dots correspond to the mean values and the numerical values plotted over each box represent the median.

tifying the main parameters which directly inform the drug safety classification, namely (I_{Kr} , I_{CaL} , I_{NaL}). These results are consistent with the importance of the main outward current I_{Kr} to assure cardiac repolarization and the counteracting effects of inward currents I_{CaL} and I_{NaL} to maintain repolarization reserve and prevent proarrhythmic events.

Table 1 summarizes the drug safety classification performance of our causally-based method compared to some state-of-the-art methods [9, 12]. Our ion-combination selection shows the best mean AUC and specificity scores, i.e. 0.94 and 94.7% respectively, outperforming the other methods. Unlike cardiac simulation classifiers, our approach does not require any cardiac action potential model to assess drug TdP risk: it only depends on the pore block model based on IC_{50} and Hill coefficients for various ion channels to calculate drug blockade. These results emphasize the importance of considering ion channels I_{CaL} and I_{NaL} with I_{Kr} to improve proarrhythmic risk classification.

For completeness’ sake, we have tested the efficacy of our method for a 4-class classification (see Sec. 2.1), which reveals that (I_{Kr} , I_{CaL}) is preferable for this task, along with a great heterogeneity across classes: this deserves further analysis, and is left as future work.

Table 1: Mean values of the performance (AUC, accuracy, sensitivity, and specificity) of our causality-based classification approach compared to some state-of-the-art methods using a set of ion currents for TdP-risk classification.

Ion combination	AUC	Acc.	Sens.	Spec.	Ref.
Kr, CaL, Na	0.91	90.9%	88%	87%	[9]
Kr, CaL, Na	NA	87%	73%	89%	[12]
Kr, CaL, NaL	0.94	93.59%	92%	94.7%	Our method

4. Conclusion

The study presented here highlights the importance of using causal discovery methods to infer ion channels selection for the drug-induced TdP risk. The obtained results suggest that I_{Kr} , I_{CaL} and I_{NaL} are the most critical ion currents to be considered for TdP risk assessment. These results are consistent with previous works (e.g., [10]) where the authors showed the crucial role of these ionic currents in the computation of *in-silico* arrhythmogenic biomarkers proposed for TdP risk assessment. In addition, the proposed causal approach can bring valuable insights concerning the downstream effects of perturbation of other ionic currents, and the obtained causal graph can further be deployed to infer the safety of new compounds. The current study has the potential to expand its scope by incorporating additional biological variables such as action potential [16], or *in-silico* biomarkers proposed in the literature [10]. This would significantly aid in selecting parameters prior to simulations, for a reduced computational time and a more reliable and rational variable selection.

Acknowledgements

This work was supported by the European Union’s Horizon 2020 Research and Innovation Program SimCardioTest project which has received funding from the European Union’s Horizon 2020 research and innovation program under grant agreement No. 101016496, and by the French government, through the 3IA Côte d’Azur Investments in the Future project managed by the National Research Agency (ANR) with the reference number ANR-19-P3IA-0002.

References

- [1] Chen W, Cai R, Zhang K, Hao Z. Causal discovery in linear non-gaussian acyclic model with multiple latent confounders. *IEEE Transactions on Neural Networks and Learning Systems* 2021;33(7):2816–2827.
- [2] Shimizu S, Hoyer PO, Hyvärinen A, Kerminen A, Jordan M. A linear non-gaussian acyclic model for causal discovery. *Journal of Machine Learning Research* 2006;7(10).
- [3] Shimizu S, Inazumi T, Sogawa Y, Hyvarinen A, Kawahara Y, Washio T, Hoyer PO, Bollen K, Hoyer P. Directlingam: A direct method for learning a linear non-gaussian structural equation model. *Journal of Machine Learning Research JMLR* 2011;12(Apr):1225–1248.
- [4] Stone JV. Independent component analysis: an introduction. *Trends in Cognitive Sciences* 2002;6(2):59–64.
- [5] Nerbonne JM, Kass RS. Molecular physiology of cardiac repolarization. *Physiological Reviews* 2005;85(4):1205–1253.
- [6] Mitchel HJ. Drug-induced hERG block and long QT syndrome. *Cardiovascular Therapeutics* 2011;29(4):251–259.
- [7] Guideline IHT. The non-clinical evaluation of the potential for delayed ventricular repolarization (QT interval prolongation) by human pharmaceuticals. S7B http://www.ich.org/products/guidelines/safety/articles/safety_guidelines.html 2005;.
- [8] Li Z, Ridder BJ, Han X, Wu WW, Sheng J, Tran PN, Wu M, Randolph A, Johnstone RH, Mirams GR, et al. Assessment of an *in silico* mechanistic model for proarrhythmia risk prediction under the CiPA initiative. *Clinical Pharmacology Therapeutics* 2019;105(2):466–475.
- [9] Kramer J, Obejero-Paz CA, Myatt G, Kuryshv YA, Bruening-Wright A, Verducci JS, Brown AM. Mice models: superior to the hERG model in predicting torsades de pointes. *Scientific Reports* 2013;3(1):1–7.
- [10] Llopis-Lorente J, Gomis-Tena J, Cano J, Romero L, Saiz J, Trenor B. *In silico* classifiers for the assessment of drug proarrhythmicity. *Journal of Chemical Information and Modeling* 2020;60(10):5172–5187.
- [11] Mistry HB. Complex versus simple models: ion-channel cardiac toxicity prediction. *PeerJ* 2018;6:e4352.
- [12] Zhou X, Qu Y, Passini E, Bueno-Orovio A, Liu Y, Vargas HM, Rodriguez B. Blinded *in silico* drug trial reveals the minimum set of ion channels for torsades de pointes risk assessment. *Frontiers in Pharmacology* 2020;1643.
- [13] Nogueira AR, Pugnana A, Ruggieri S, Pedreschi D, Gama J. Methods and tools for causal discovery and causal inference. *Wiley Interdisciplinary Reviews Data Mining and Knowledge Discovery* 2022;12(2):e1449.
- [14] Foraita R, Spallek J, Zeeb H. *Directed Acyclic Graphs*. New York, NY: Springer New York, 2014; 1481–1517.
- [15] Ruta D, Gabrys B. Classifier selection for majority voting. *Information Fusion* 2005;6(1):63–81. ISSN 1566-2535. *Diversity in Multiple Classifier Systems*.
- [16] Jeong DU, Yoo Y, Marcellinus A, Lim KM. Application of convolutional neural networks using action potential shape for *in-silico* proarrhythmic risk assessment. *Biomedicine* 2023;11(2):406.

Address for correspondence:

Safaa Al-Ali
 Centre INRIA d’Université Côte d’Azur, EPIONE team,
 06902 Sophia Antipolis, France.
 safaa.al-ali@inria.fr