

Biatrial Modelling for In Silico Prediction of Atrial Fibrillation Inducibility

Semhar B Misghina¹, Jose A Solis-Lemus², Edward J Vigmond³, Steven A Niederer², Caroline H Roney¹

¹Queen Mary University of London, London, United Kingdom

²Imperial College London, London, United Kingdom

³University of Bordeaux, Bordeaux, France

Abstract

Atrial fibrillation (AF) is a cardiac disorder characterised by rapid atrial contractions. Current treatments, including ablation, vary in effectiveness. Recent mechanistic modelling studies have highlighted the significance of the right atrium (RA) in predicting AF outcomes, although its role remains unclear. This study employs a novel open-source biatrial modelling pipeline to assess AF inducibility and monitor AF dynamics on clinical timescales.

Patient-specific models were created from late gadolinium enhancement MRI (LGE-MRI) scans of 20 patients. Manual RA and left atrial (LA) segmentation, fibrosis mapping in pre-processing, and calculation of atrial coordinates to incorporate atrial structures and fibres were performed. These personalised models were simulated and post-processed to assess the AF wavefront patterns.

RA integration significantly increased rotor activity and total phase singularities (PS) within the LA posterior walls and reduced conduction velocity, indicating greater potential for AF sustainability. LA exhibited a higher mean PS density (3.8 rotors/cm²) than RA (2.1 rotors/cm²), indicating regions prone to re-entry or wavefront break-up.

The modelling pipeline highlights the potential of biatrial models to efficiently predict AF outcomes, enabling personalised therapies and comparisons of ablation approaches and anti-arrhythmic drug therapies.

1. Introduction

Atrial fibrillation (AF) is a prevalent cardiac arrhythmia characterised by abnormal electrical impulses [1]. Current anti-arrhythmic drugs are sub-optimal, making radio-frequency catheter ablation (RFCA) the gold standard [1]. RFCA poses challenges in predicting ablation targets, leading to the adoption of LGE-MRI for 3D mapping and optimised therapy. These imaging advances enhance personalised therapy and streamline clinical workflows [2]. Prior research has primarily focused on LA

modelling to understand AF; however, the potential of biatrial models in predicting AF inducibility remains under-explored [3,4]. The RA presents challenges due to limited pre-labelled datasets and geometric variability in manual segmentation [2]. This study addresses these constraints as cardiac modelling advances towards large-scale in silico studies and personalised predictions, which holds potential for patient-specific and population digital twins [3,4].

We present a biatrial modelling pipeline using LGE-MRI data from 20 patients. To address RA segmentation challenges, we employed segmentation tools with an MRI atlas. Patient-specific bilayer meshes incorporated atrial fibres and structures from a fibre atlas using atrial coordinates. We conducted finite element simulations and evaluated PS density maps for AF inducibility assessment.

2. Materials and Methods

2.1. Patient Cohort

This study used LGE-MRI scans of the atria from 20 patients in the Atrial Segmentation Challenge Dataset (2018) [5]. These scans had a spatial resolution of 0.625 x 0.625 x 0.625 mm³ with variable spatial dimensions (576 x 576 x 88 or 640 x 640 x 88 pixels). Ethical guidelines were followed, and all patients provided informed consent [6].

2.2. Image Segmentation

Atrial anatomical models were created from LGE-MRI data using 3D Slicer software (version 5.3.0). This process involved importing DICOM data, manually tracing LA and RA from individual 2D cross-sectional images, and exporting the 3D models in NIfTI format for mesh pre-processing. To ensure accurate labelling, pre-labelled LA datasets and a reference MRI atlas were employed and models were validated by a cardiac modelling expert. Segmented meshes were saved in the VTK file format for all cases. All the steps in the model construction process are shown in Figure 1.

3.2. Fibrotic Remodelling

Fibrotic modelling, based on LGE intensities, revealed significant diversity among the 20 anatomies, as shown in Figure 3A. Models that required corrections in clipping and segmentation were revised, and an example clipping is shown in Figure 3B. The diverse atrial geometry posed challenges in landmark selection, but expert validation ensured accuracy. The meshes were combined to create biatrial models, which were validated visually and by fibre mapping, with two excluded due to insufficient segmentation.




Figure 3. **LGE intensities projected on the 3D LA and RA meshes to incorporate fibrosis.** (A) Fibrosis mapping showing healthy (blue) and scarred tissue (red) in the AP, posteroanterior (PA), and right atrial oblique (RAO) views. The colour axis indicates the IIR values. (B) Clipped LA and RA surfaces at PV, CS, VC, TV, and MV.

3.3. Fibre Modelling

Our approach, similar to Roney et al.(2022) methods, maps atrial structures and fibres in both the LA and RA [3]. Figure 4 shows fibre configurations in the biatrial models, highlighting their significance in personalisation. The exclusion of complex atrial meshes streamlined fibre integration, facilitating biophysical modelling.




Figure 4. **Atrial fibres from the Labarthe et al.(2014) atlas were modelled using 2D UAC systems for (1) LA and (2) RA.** (B) Epicardial and (C) endocardial surfaces are depicted in AP and axial views as streamlines [2].

3.4. Biophysical Models

Finite element simulations assessed the AF dynamics over 15s in a biatrial model. Figure 5 shows slower conduction in the LA posterior walls (LAPW), indicating significantly slower mean conduction velocity in the LA (31.7 cm/s) than in the RA (48.2 cm/s).




Figure 5. **Transmembrane potential maps from AF simulations using openCARP software depict AF induction at 0ms, 100ms, 200ms, and 300ms.** Arrows indicate wavefront patterns in (A) AP and (B) PA views, with the colour bar showing the transmembrane potential in mV.

Figure 6 shows PS density maps with increased rotor activity and PS hotspots in the LAPW, suggesting re-entry prone areas and a higher likelihood of sustained AF. The LA exhibits a maximal phase singularity density of 3.8 rotors/cm², while the RA has 2.1 rotors/cm².




Figure 6. **Post-processed PS density maps from AF simulations are shown in AP, PA, axial, and RAO views.** These maps show PS hotspot localisation in the LAPW.

4. Discussion

In this study, we developed biatrial models using LGE-MRI data to predict AF inducibility in 20 patients. Rotors, functional re-entry mechanisms in AF, are influenced by ionic variations and scarring. Integrating RA significantly affected AF dynamics, with PS density maps indicating increased rotor activities in the LAPW, which are markers of AF recurrence. These rotor locations aligned with regions of scar tissue identified during mesh pre-processing.

Our biatrial models align with the findings of Calo et al.(2006), highlighting the key role of RA re-entrant arrhythmia in AF development and ablation treatment. This study also identified conduction issues and chaotic impulses in certain atrial walls [11]. Our work extends the study of Nagel et al.(2021), which focused on biatrial statistical modelling and imaging metrics for assessing AF

activity. Our study assesses AF dynamics using patient-specific biatrial models and expands the literature on the role of RA in AF recurrence, complementing Hopman et al.(2023) work on RA fibrotic remodelling and ablation approaches [4, 12]. This study assessed the specific effects of RA and LA on AF recurrence, potentially informing personalised catheter ablation strategies.

The biatrial modelling framework has limitations, particularly in manual segmentation which introduces intra-operator variability. Although developing an autonomous biatrial modelling pipeline is not the focus of this study, it could minimise errors and enhance region assignment repeatability [13]. Although model personalisation is important in identifying AF trends, our biatrial models do not account for specific patient attributes such as co-morbidities, limiting their applicability in diverse populations [3].

This study aimed to predict AF inducibility using biatrial models. It also offers the potential for assessing ablation strategies to treat AF recurrence. Future work will extend this work to the UK Biobank longitudinal dataset, enabling long-term post-AF treatment outcome prediction. Our objectives will address the challenges of inferring missing atrial longitudinal data, constructing models from low-resolution imaging data, and integrating ECG data using biophysical models and machine learning. These research directions advance our understanding of AF mechanisms and aim to streamline clinical workflows.

5. Conclusion

Our study introduces a biatrial model cohort to assess AF patterns and highlights the role of the RA in predicting AF inducibility. These findings highlight the potential of biatrial substrates in the development of personalised patient-specific and population digital twins.

Acknowledgments

This research was funded by a UKRI Future Leaders Fellowship (MR/W004720/1).

References

- [1] Pashakhanloo F. High-Resolution Whole-Heart Imaging and Modeling for Studying Cardiac Arrhythmia. Ph.D. thesis, Johns Hopkins University, 2016.
- [2] Zhuang X. Automatic Whole Heart Segmentation Based on Image Registration. Ph.D. thesis, UCL (University College London), 2010.
- [3] Roney CH, Sim I, Yu J, Beach M, Mehta A, Solis-Lemus JA, Kotadia I, Whitaker J, Corrado C, Razeghi O, et al. Predicting Atrial Fibrillation Recurrence by Combining Population Data and Virtual Cohorts of Patient-Specific Left Atrial Models. *Circulation Arrhythmia and Electrophysiology* 2022;15(2):e010253.
- [4] Hopman LH, Visch JE, Bhagirath P, van der Laan AM, Mulder MJ, Razeghi O, Kemme MJ, Niederer SA, Allaart CP, Götte MJ. Right Atrial Function and Fibrosis in Relation to Successful Atrial Fibrillation Ablation. *European Heart Journal Cardiovascular Imaging* 2023;24(3):336–345.
- [5] Borra D, Andalò A, Paci M, Fabbri C, Corsi C. A Fully Automated Left Atrium Segmentation Approach from Late Gadolinium Enhanced Magnetic Resonance Imaging Based on a Convolutional Neural Network. *Quantitative Imaging in Medicine and Surgery* 2020;10(10):1894.
- [6] McGann C, Kholmovski E, Blauer J, Vijayakumar S, Haslam T, Cates J, DiBella E, Burgon N, Wilson B, Alexander A, et al. Dark Regions of No-Reflow on Late Gadolinium Enhancement Magnetic Resonance Imaging Result in Scar Formation after Atrial Fibrillation Ablation. *Journal of the American College of Cardiology* 2011;58(2):177–185.
- [7] Razeghi O, Solís-Lemus JA, Lee AW, Karim R, Corrado C, Roney CH, de Vecchi A, Niederer SA. Cemrgapp: An Interactive Medical Imaging Application with Image Processing, Computer Vision, and Machine Learning Toolkits for Cardiovascular Research. *SoftwareX* 2020;12:100570.
- [8] Gunturiz-Beltrán C, Borràs R, Alarcón F, Garre P, i Ventura RMF, Benito EM, Caixal G, Althoff TF, Tolosana JM, Arbelo E, et al. Quantification of Right Atrial Fibrosis by Cardiac Magnetic Resonance: Verification of the Method to Standardize Thresholds. *Revista Espanola de Cardiologia English Edition* 2023;76(3):173–182.
- [9] Roney CH, Pashaei A, Meo M, Dubois R, Boyle PM, Trayanova NA, Cochet H, Niederer SA, Vigmond EJ. Universal Atrial Coordinates applied to Visualisation, Registration and Construction of Patient Specific Meshes. *Medical Image Analysis* 2019;55:65–75.
- [10] Courtemanche M, Ramirez RJ, Nattel S. Ionic Targets for Drug Therapy and Atrial Fibrillation-Induced Electrical Remodeling: Insights from a Mathematical Model. *Cardiovascular Research* 1999;42(2):477–489.
- [11] Calò L, Lamberti F, Loricchio ML, De Ruvo E, Colivicchi F, Bianconi L, Pandozi C, Santini M. Left Atrial Ablation versus Biatrial Ablation for Persistent and Permanent Atrial Fibrillation: A Prospective and Randomized Study. *Journal of the American College of Cardiology* 2006;47(12):2504.
- [12] Nagel C, Schuler S, Dössel O, Loewe A. A Bi-Atrial Statistical Shape Model for Large-Scale In Silico Studies of Human Atria: Model Development and Application to ECG Simulations. *Medical Image Analysis* 2021;74:102210.
- [13] Aguado AM, Olivares AL, Yagüe C, Silva E, Nuñez-García M, Fernandez-Quilez Á, Mill J, Genua I, Arzamendi D, De Potter T, et al. In Silico Optimization of Left Atrial Appendage Occluder Implantation using Interactive and Modeling Tools. *Frontiers in Physiology* 2019;10:237.

Address for correspondence:

Semhar Misghina
Queen Mary University of London, UK
s.b.misghina@qmul.ac.uk